

Références de l'article de Joanie Belleau intitulé : Étude de l'expérience de femmes consultant à l'urgence gynécologique du CHU Sainte-Justine pour un avortement spontané durant le 1er trimestre de grossesse ne nécessitant pas une intervention chirurgicale

- CHU Sainte-Justine. (2011). *À propos de nous, Pôles d'excellence*, récupéré le 4 février 2013 http://www.chu-sainte-justine.org/Apropos/page.aspx?ID_Menu=617&ID_Page=4739&ItemID=2a
- CHU Sainte-Justine. (2013). *Rapport des activités du triage gynéco-obstétrical*, STAT-DEV, Montréal.
- Everett, C. (1997). Incidence and outcome of bleeding before the 20th week of pregnancy: prospective study from general practice. *BMJ: British Medical Journal (International Edition)*, 315(7099), 32-34.
- Gerber-Epstein, P., Leichtentritt, R. D., & Benyamini, Y. (2009). The experience of miscarriage in first pregnancy: the women's voices. *Death Studies*, 33(1), 1-29.
- Giorgi, A. (1970). *Psychology as a human science : A phenomenologically based approach*. New York: Harper & Row. 240p.
- Huacahuari, N. (2012). Emergencies in early pregnancy. *Emergency Medicine Clinics of North America*, 30(4), 837-847.
- Loiselle, C. G., Profetto-McGrath, J., Polit, D. F., & Beck, C. T. (2007). *Méthodes de recherche en sciences infirmières*. Montréal : ERPI inc.
- Murphy, F. A., Lipp, A., & Powles, D. L. (2012). Follow-up for improving psychological well being for women after a miscarriage. *Cochrane Database of Systematic Reviews*(3).
- Rando, T. A. (1985). Bereaved parents: Particular difficulties, unique factors, and treatment issues. *Social Work*, 30(1), 19-23.
- Rogers, C. H., Floyd, F. J., Seltzer, M. M., Greenberg, J., & Hong, J. (2008). Long-term effects of the death of a child on parents' adjustment in midlife. *Journal of Family Psychology*, 22(2), 203-211.
- Tang, O. S., & Ho, P. C. (2006). The use of misoprostol for early pregnancy failure. *Current Opinion in Obstetrics & Gynecology*, 18(6), 581-586.